

A MEALY MACHINE WITH POLYNOMIAL GROWTH OF IRRATIONAL DEGREE

LAURENT BARTHOLDI AND ILLYA I. REZNYKOV

ABSTRACT. We consider a very simple Mealy machine (two non-trivial states over a two-symbol alphabet), and derive some properties of the semigroup it generates. It is an infinite, finitely generated semigroup, and we show that the growth function of its balls behaves asymptotically like n^α , for $\alpha = 1 + \log 2 / \log \frac{1+\sqrt{5}}{2}$; that the semigroup satisfies the identity $g^6 = g^4$; and that its lattice of two-sided ideals is a chain.

1. INTRODUCTION

Algebraic objects may be defined by universal properties, like the identities they satisfy, or a presentation as a quotient of two free objects. They may also be defined by their action on simpler objects, viz, for an algebra, as endomorphisms of a vector space; for a group, as permutations of a set, and for a semigroup, as mappings of a set.


If the semigroup to be defined is infinite, it naturally must act on an infinite set; extra conditions must be imposed on the action to ensure that it remains describable (by a finite sequence of mathematical symbols, say). A specially interesting class of semigroups appear by requesting that the infinite set be the set X^* of words over a finite alphabet, and that the action be given by a finite-state automaton.

The growth function $\gamma(\ell)$ of a finitely generated semigroup Γ — the number of semigroup elements that can be obtained as products of at most ℓ generators — is an important invariant of the semigroup. It depends on the chosen generating set, but its asymptotics do not. This function is at most exponential. If it is bounded by a polynomial, then Γ is of *polynomial growth*; following [6], its *Gelfand-Kirillov dimension* (abbreviated GK dimension) is then defined as the infimum of those α such that $\gamma(\ell)/\ell^\alpha$ does not converge to 0.

The “Bergman gap” [8, Theorem 2.5] asserts that a semigroup may have GK dimension 0, 1, or ≥ 2 ; by a result by Warfield [11], there exist semigroups of GK dimension β for any $\beta \geq 2$. Belov and Ivanov [3, 4] construct finitely presented semigroups with non-integer GK dimension. Shneerson [10] constructs relatively free (i.e., free relative to an identity) semigroups of intermediate growth (asymptotically $m^{\log m}$). His proof involves Fibonacci numbers, as does the present construction.

This paper describes the semigroup $\Gamma(I)$ generated by the 3-state automaton I in Figure 1 (see §3 for the definition of a semigroup generated by an automaton), and proves that its GK dimension is irrational. A much more precise statement appears in Theorem 2.4.

The main purpose of this paper is to show that a very simple-minded finite-state automaton can produce a semigroup with a highly unusual growth pattern,

FIGURE 1. The automaton I

asymptotically n^α for an irrational α . Other exotic types of growth of automata have also been discovered; for example, of type $e^{\sqrt{n}}$ in [2], and of type $n^{\log(n)/2\log m}$ for integer m , in [9]. It is intriguing that in all these cases the determination of the growth function relies on enumerations of partitions with certain constraints.

1.1. Plan. The next section gives an automata-free version of the semigroup studied in this paper, along with a presentation of its main results. Section 3 gives the necessary definitions of semigroups generated by automata. Statements whose proof are too long to fit smoothly in the text are stated as underlined. Their proofs appear in Section 5.

1.2. Notation. All actions are written on the right in this paper. The identity is written e . We use $=$ for equality of group elements, and $\equiv$ for graphical (letter-by-letter) equality of words representing these elements. We denote by X^* the free monoid on the set X . The length of a word w is denoted by $\|w\|$; this is also the length of a minimal word representing a semigroup element.

The integers are written $\mathbb{Z}$, and the naturals (containing 0) are written $\mathbb{N}$. Congruence is written $\equiv$, and the ‘mod 2’ operation (remainder after division by 2) is written $n\%2$ as in the C programming language.

Sequences of the form $f_a f_{a+2} \dots f_b$, sometimes simply written $f_a \dots f_b$, appear throughout the paper. They are taken to be e if $a > b$, and $f_a f_{a+2} f_{a+4} \dots f_{b-2} f_b$ otherwise.

1.3. Acknowledgments. We are very grateful to the anonymous referee who suggested substantial improvements to the paper, in particular a clarification of the role of rewriting systems.

2. MAIN RESULTS

Consider the following transformations s, f of the integers $\mathbb{Z}$:

$$(1) \quad \begin{aligned} x^s &= \begin{cases} x-1 & \text{if } x \text{ is odd,} \\ x+1 & \text{if } x \text{ is even;} \end{cases} \\ x^f &= \begin{cases} x-2^n-1 & \text{if there exists } n \geq 0 \text{ such that } x \equiv 3 \cdot 2^n \pmod{2^{n+2}}, \\ x+3 \cdot 2^n-1 & \text{if there exists } n \geq 0 \text{ such that } x \equiv 2^n \pmod{2^{n+2}}, \\ -1 & \text{if } x = 0. \end{cases} \end{aligned}$$

Note that f is uniquely defined, because every $x \in \mathbb{Z}$ is either 0 or of the form $2^n y$ for unique $n \geq 0$ and odd y , with either $y \equiv 1$ or $y \equiv 3$ modulo 4.

2.1. The semigroup. Let F be the semigroup generated by the transformations s and f . Define furthermore the elements f_n of F by $f_1 = s$, $f_2 = f$, and inductively $f_n = f_{n-2}f_{n-1}$ for $n \geq 3$. For example, $f_3 = sf$, $f_4 = fsf$, $f_5 = sf^2sf$. These words are sometimes called the “Fibonacci sequence” — see §2.2 for the connection.

Recall [5] that a rewriting system for a semigroup Γ generated by a set Q is a set of equations, called *rules*, of the form $\ell \rightarrow r$, with $\ell, r \in Q^*$. An elementary reduction in a word $w \in Q^*$ is the replacement of a subword equal to the left-hand side of a rule by the right-hand side; if no elementary reduction is possible, the word is *reduced*. A rewriting system is *terminating* if there is no $w \in Q^*$ to which an infinite sequence of elementary reductions can be applied; and it is *confluent* if the reduced word obtained after applying as many elementary reductions as possible does not depend on the choice of elementary reductions. It is *complete* if it is terminating and confluent; the set of reduced words is then in bijection with the semigroup, through the natural evaluation map $Q^* \rightarrow \Gamma$, and is called a *normal form* for Γ .

For $n \geq 1$, define words r_n and r'_n over the alphabet $\{s, f\}$ as follows:

$$\begin{aligned} (2) \quad & r_1 = s^2, \quad r'_1 = e, \\ & r_n = f_{n+1}f_n^2, \\ (3) \quad & r'_n = f_{n\%2+1}(f_{n\%2+5}f_{n\%2+7} \cdots f_{n-1}f_{n+1})f_n. \end{aligned}$$

In particular, $r_2 = sf^3$ and $r'_2 = sf$, and $r_3 = f_4f_3^2$ and $r'_3 = f_4$. For $n \geq 3$, we have $\|r'_n\| = \|r\| - 4$.

Theorem 2.1. *The semigroup F is infinite, and admits as presentation*

$$\langle s, f \mid r_n = r'_n \text{ for all } n \geq 1 \rangle,$$

Furthermore, after the relations $sw = sw'$ (which occur for even n) are replaced by the equivalent relations $w = w'$, this presentation is a complete rewriting system.

2.2. A normal form. Let Φ_n denote the sequence of Fibonacci numbers, defined by $\Phi_1 = \Phi_2 = 1$ and $\Phi_n = \Phi_{n-2} + \Phi_{n-1}$ for $n \geq 3$. Then

Theorem 2.2. *Every element g of F admits a unique representation as a word of the form*

$$(4) \quad w_g = s^\epsilon f_{i_1} f_{i_2} \cdots f_{i_m} \cdots f_{i_n},$$

for some $n \geq 0$, some $\epsilon \in \{0, 1\}$, and some indices $i_1, \dots, i_n$ satisfying

$$3 \leq i_1, i_1 + 1 < i_2, \dots, i_{m-1} + 1 < i_m > i_{m+1} > \cdots > i_n \geq 1.$$

We call i_m the maximal index of w_g .

When spelled out in the generating set $\{s, f\}$, this representation of g is essentially minimal: if i_1 is even, then w_g is the unique minimal-length representation of g , while if i_1 is odd, then an initial s^2 must be cancelled to obtain the unique minimal representation of g . Its length (see §3.3) is

$$\|g\| = (-1)^{i_1} \epsilon + \Phi_{i_1} + \cdots + \Phi_{i_n}.$$

By a slight extension of the definition of rewriting system, let us admit rules of the form $\wedge \ell \rightarrow r$ that mean that the subword ℓ may be replaced in w by the word r if it is a prefix of w . We will actually consider the following rewriting system; it is on an infinite generating set, but its rules are much simpler, since their left-hand sides have length at most 3.

Theorem 2.3. *On the generating set $\{f_i : i \geq 1\}$, the semigroup F admits a complete rewriting system with rules*

- (N1) $f_1^2 \longrightarrow e$
- (N2) $f_a f_{a+1} \longrightarrow f_{a+2} \quad (a \geq 1)$
- (N3) $f_a^2 \longrightarrow f_{a-2} f_{a+1} \quad (a \geq 3)$
- (N4) $f_a f_2^2 \longrightarrow f_a \quad (a \geq 2)$
- (N5) $\wedge f_2 \longrightarrow f_1 f_3$
- (N6) $f_{a+1} f_a f_{a+3} \longrightarrow f_{a+3} f_{a+2} \quad (a \geq 1)$
- (N7) $f_{a+2} f_a f_{a+3} \longrightarrow f_a f_{a+3} f_{a+2} \quad (a \geq 1)$
- (N8) $f_{a+p} f_a f_{a+q} \longrightarrow \underbrace{f_{a+p-2} \cdots f_a}_{p-1 \text{ terms}} f_2^2 \cdots f_{a-2}^2 f_{a-1} f_{a+q}$
 $(a \geq 2, p \geq 1, q \geq 2 \text{ even})$
- (N9) $f_{a+p} f_a f_{a+q} \longrightarrow \underbrace{f_{a+p-2} \cdots f_{a+2}}_{p-3 \text{ terms}} f_2^2 \cdots f_a^2 f_{a+q} \quad (a \geq 1, p \geq 3, q \geq 3 \text{ odd})$
- (N10) $f_{1+p} f_1 f_{1+q} \longrightarrow f_{1+p} f_2 f_4 \cdots f_q \quad (p \geq 1, q \geq 2 \text{ even}).$

From now on, we identify g with w_g , and call the word w_g the *normal form* of g . We will never use a different notation for a semigroup element and its normal form.

2.3. Growth. Using this minimal representation, the (ball) growth function of F ,

$$\gamma(\ell) = \#\{g \in F : \|g\| \leq \ell\},$$

may be quite precisely estimated; namely

Theorem 2.4. *There are constants $C, D > 0$ such that the growth function of F satisfies*

$$C\ell^\alpha \leq \gamma(\ell) \leq D\ell^\alpha$$

for all $\ell \in \mathbb{N}$, where $\varphi = (1 + \sqrt{5})/2$, and $\alpha = 1 + \log 2 / \log \varphi \approx 2.4401$, and $C = \frac{2\sqrt{5}^\alpha}{\sqrt{5}\varphi^2(2\varphi-1)(2\varphi)^\alpha}$, and $D = \frac{2\sqrt{5}^\alpha}{\sqrt{5}\varphi^2(2\varphi-1)}$.

Therefore, the Gelfand-Kirillov dimension of F is α .

Experimental computations indicate that actually the function $\gamma(\ell)/\ell^\alpha$ does not converge, but oscillates between ≈ 0.201 and ≈ 0.205 , reaching its maxima at Fibonacci numbers.

2.4. Identities. We have not determined the complete set of identities satisfied by F ; nevertheless, we know

Theorem 2.5. *The semigroup F satisfies the identity $g^6 = g^4$.*

2.5. The ideal structure of F . We describe in this subsection the quotient and ideal structure of F . First, we may consider for all $n \in \mathbb{N}$ the quotient of F , denoted W_n , acting as transformations on X^n . Let us denote, for $n \in \mathbb{N}$,

$$z_n = \begin{cases} f_3 f_5 \cdots f_{n+2} & \text{if } n \text{ is odd,} \\ f_4 f_6 \cdots f_{n+2} & \text{if } n \text{ is even.} \end{cases}$$

Then we have

Theorem 2.6. (1) *The semigroup W_n is presented as follows:*

- $W_n = \langle s, f \mid r_k = r'_k, 1 \leq k \leq n+2, \text{ and } sz_n = fz_n = z_n, f_{n+2}f_{n+1} = z_n \rangle.$
- (2) *The elements of W_n may be described as all normal forms $f_1 \cdots f_{i_m} \cdots f_{i_t}$ whose maximal index i_m is $< n+2$, normal forms of the form $z_n f_{i_{m-1}} \cdots f_{i_t}$ for $n \geq i_{m-1} > \cdots > i_t$, and those normal forms of maximal index $n+2$ that include neither z_n nor $f_{n+2}f_{n+1}$.*
- (3) *The semigroup W_n has order $\sum_{k=1}^n 2^{k+2}\Phi_k - 2^n - 2^{n+1}\Phi_n + 2$.*

We identify for the remainder of this section the quotient semigroup W_n with the set of normal forms of its elements, which then form a subset of F .

Corollary 2.7. *The Hausdorff dimension (in the sense of §3.6) of F is 0.*

Proof. Using the count of elements in W_n , we have

$$\#W_n \leq 2 + \sum_{k=1}^n 2^{k+2}\Phi_k \leq \frac{4}{\sqrt{5}} \frac{(2\varphi)^{n+1} - 1}{2\varphi - 1} \leq (2\varphi)^{n+1},$$

so

$$\text{Hdim}(F) = \liminf_{n \rightarrow \infty} \frac{\log \#W_n}{\log \# \text{End}(X^n)} = \liminf_{n \rightarrow \infty} \frac{\log(2\varphi)^{n+1}}{\log 2^{2(2^n-1)}} = 0. \quad \square$$

Corollary 2.8. *Let g be an arbitrary element of F with maximal index $n \geq 3$. Then its trace (in the sense of §3.5) satisfies*

$$\tau(g) = \begin{cases} 2^{2-n} & \text{if } g \text{ includes } z_{n-2} \text{ or } f_n f_{n-1} \text{ in its normal form,} \\ 2^{3-n} & \text{otherwise.} \end{cases}$$

The ideal $I_n = Ff_{n+3}F$ coincides with F_{2-n} , as defined in §3.5.

Theorem 2.9. *All two-sided ideals of F are of the form Ff_nF , and equivalently of the form F_ξ as defined in §3.5. They therefore form a unique chain.*

Recall that if I is an ideal of F , the quotient F/I is the semigroup whose elements are the equivalence classes of the congruence $I \times I \cup \Delta$, where Δ is the identity relation (= diagonal) of F .

Proposition 2.10. *For $n \geq 2$, the quotient F/I_n has order $\sum_{k=1}^n 2^{k+2}\Phi_k - 2^{n+1} - 2^{n+1}\Phi_n + 3$. It is obtained from W_n by identifying together all idempotents of rank 1, i.e. all maps $f_p : X^n \rightarrow X^n$ defined by $f_p(x_1 \dots x_n) = p$, for all $p \in X^n$.*

Proof. By Theorem 2.6, the quotient F/I_n is a quotient of W_n . The transformation f_{n+3} induces a transformation of rank 1 on X^n , and its left multiples, which are identified in F/I_n , are the maps f_p in the statement of the Proposition. $\square$

3. DEFINITIONS

A *finite-state automaton*, also called *Mealy machine*, is comprised of the following data: a set X , the *alphabet*; a set Q , the *states*; a function $\tau : X \times Q \rightarrow Q$, the *transition*; a function $\pi : X \times Q \rightarrow X$, the *output*. In this section we suppose that the alphabet is $X = \{0, \dots, d\}$; we will later specialize to $d = 1$.

Such an automaton $\mathcal{A}$ is usually represented, as in the figure above, as a graph. The states are represented by vertices, and there is an edge from vertex q to vertex r , labelled $i \rightarrow o$, whenever $\tau(i, q) = r$ and $\pi(i, q) = o$.

3.1. Action. States $q \in Q$ of $\mathcal{A}$ yield transformations of X^* , which are defined simultaneously as follows:

$$(5) \quad (i_1 \dots i_n)^q = \pi(i_1, q)(i_2 \dots i_n)^{\tau(i_1, q)}.$$

These transformations generate a semigroup $\Gamma(\mathcal{A}) = \langle Q \rangle$, called the *semigroup generated by $\mathcal{A}$* .

3.2. Decomposition. Yet another way of describing the automaton $\mathcal{A}$ is via its *semigroup decomposition*. Given $g \in \Gamma$, it yields by (5) a transformation π_g of X by restriction to length-1 words; and, for all $i \in X$, a transformation $iX^* \rightarrow i^{\pi_g}X^*$, again by restriction. By (5), the composition $X^* \rightarrow iX^* \rightarrow i^{\pi_g}X^* \rightarrow X^*$ is again an element of Γ , which we denote by g_i . We write

$$(6) \quad \phi(g) = \llbracket g_0, \dots, g_d \rrbracket \pi_g$$

for the decomposition of $g \in \Gamma$, with $g_i \in \Gamma$ and $\pi_g : X \rightarrow X$. Multiplication of such decompositions obeys the rule

$$(7) \quad \llbracket g_0, \dots, g_d \rrbracket \pi_g \llbracket h_0, \dots, h_d \rrbracket \pi_h = \llbracket g_0 h_{\pi_g(0)}, \dots, g_d h_{\pi_g(d)} \rrbracket \pi_g \pi_h.$$

It is therefore sufficient to know the decomposition of generators, and these are determined by the transition and output functions $\tau, \pi : X \times Q \rightarrow Q$ of $\mathcal{A}$, by

$$\phi(q) = \llbracket \tau(0, q), \dots, \tau(d, q) \rrbracket (i \mapsto \pi(i, q) \forall i).$$

3.3. Metrics. On the semigroup Γ generated by a set Q , define the “norm”

$$\|g\| = \min\{\ell \in \mathbb{N} : g = q_1 \cdots q_\ell, \quad q_i \in Q \forall i\}.$$

(we have $\|gh\| \leq \|g\| + \|h\|$, which justifies calling it a norm.) The *ball growth function* is the function $\gamma : \mathbb{N} \rightarrow \mathbb{N}$ defined by

$$\gamma(\ell) = \#\{g \in \Gamma : \|g\| \leq \ell\};$$

it measures the volume growth of balls in the discrete normed space Γ .

3.4. Contraction. An automaton is *contracting* [7] if there are constants C and $\eta < 1$ such that whenever $\phi(g) = \llbracket g_0, \dots, g_d \rrbracket \pi_g$, we have

$$\|g_i\| \leq \eta \|g\| + C.$$

Contracting automata are most studied, in part because contraction gives a natural strategy of proof by induction.

3.5. Traces. Let g be a transformation of X^* given by an automaton. Then for all $n \in \mathbb{N}$ we have $\#g(X^{n+1}) \leq \#X \#g(X^n)$, so the limit

$$\tau(g) := \lim_{n \rightarrow \infty} \frac{\#g(X^n)}{\#X^n} \in [0, 1]$$

exists. We call it the *trace* of g . The following facts are easily checked:

- $\tau(gh) \leq \min\{\tau(g), \tau(h)\}$;
- $\tau(g) = 1$ if and only if g is invertible.

Therefore, if Γ is the semigroup generated by an automaton, we may define for any $\xi \in [0, 1]$ a two-sided ideal $\Gamma_\xi = \{g \in \Gamma : \tau(g) \leq \xi\}$. We have $\Gamma_\xi = \Gamma$ if and only if $\xi = 1$.

3.6. Hausdorff dimension. Let Γ be a semigroup acting on a rooted tree X^* . There is another metric on Γ , defined by

$$d(g, h) = \exp(-\max\{n : v^g = v^h \text{ for all } v \in X^n\}),$$

with the convention that $\exp(-\infty) = 0$. This turns Γ into a metric space of diameter at most 1. The *Hausdorff dimension* of Γ is defined as the Hausdorff dimension of this metric space. Let $\text{End}(X^n)$ denote the set of prefix-preserving maps $X^n \rightarrow X^n$. If $\#X = d + 1$, then

$$\#\text{End}(X^n) = ((d + 1)^{d+1})^{((d+1)^n - 1)/d}.$$

Then the Hausdorff dimension of Γ is given by the formula

$$\text{Hdim}(\Gamma) = \liminf_{n \rightarrow \infty} \frac{\log \#\Gamma_n}{\log \#\text{End}(X^n)},$$

where Γ_n is the quotient of Γ acting as a transformation semigroup of X^n ; see [1] for an analogous definition in the case of profinite groups.

4. THE AUTOMATON I AND THE SEMIGROUP F

We now consider the automaton I from Figure 1. This automaton has three states f, s, e and a two-letter alphabet $X = \{0, 1\}$.

Consider the following transformations σ, ζ of X :

$$\sigma(i) = 1 - i, \quad \zeta(i) = 0.$$

Then the decomposition of the states are

$$\phi(f) = \ll s, f \gg \zeta, \quad \phi(s) = \ll e, e \gg \sigma, \quad \phi(e) = \ll e, e \gg.$$

Clearly e acts as the identity transformation, and s is invertible, of order 2.

4.1. Action on integers. We show now that the semigroup defined in Section 2 by its action on integers is the semigroup generated by an automaton.

Theorem 4.1. *The semigroups F and $\Gamma(I)$ are isomorphic.*

Proof. The action of $F = \langle s, f \rangle$ given in (1) extends to a continuous action of F on the 2-adics $\mathbb{Z}_2$. Consider the following bijection Θ between X^∞ and $\mathbb{Z}_2$:

$$\Theta(x_1 x_2 \dots) = \sum_{i=1}^{\infty} (1 - x_i) 2^{i-1}.$$

Let us denote temporarily $\Gamma(I) = \langle \tilde{s}, \tilde{f} \rangle$. The theorem will follow from $\Theta(x^{\tilde{s}}) = \Theta(x)^s$ and $\Theta(x^{\tilde{f}}) = \Theta(x)^f$ for all $x \in X^\infty$.

Consider therefore $x = x_1 x_2 \dots \in X^\infty$, and write $y = x_2 x_3 \dots$. If $x_1 = 0$, then

$$\Theta(x)^s = \Theta(0y)^s = (1 + 2\Theta(y))^s = 2\Theta(y) = \Theta(1y) = \Theta(x^{\tilde{s}}),$$

while if $x_1 = 1$, then

$$\Theta(x)^s = \Theta(1y)^s = (2\Theta(y))^s = 1 + 2\Theta(y) = \Theta(0y) = \Theta(x^{\tilde{s}}).$$

Let next $n \in \mathbb{N} \cup \{\infty\}$ be maximal such that $x_1 = \dots = x_n = 1$. If $n = \infty$, then

$$\Theta(x)^f = \Theta(11\dots)^f = 0^f = -1 = \Theta(00\dots) = \Theta(x^{\tilde{f}});$$

otherwise, $x = 1\dots 10x_{n+2}x_{n+3}\dots$; write $y = x_{n+3}x_{n+4}\dots$. If $x_{n+2} = 0$, then

$$\Theta(x)^f = (2^n + 2^{n+1} + 2^{n+2}\Theta(y))^f = 2^{n+1} - 1 + 2^{n+2}\Theta(y) = \Theta(0\dots 001y) = \Theta(x^{\tilde{f}}),$$

while if $x_1 = 1$, then

$$\Theta(x)^f = (2^n + 2^{n+2}\Theta(y))^f = 2^{n+2} - 1 + 2^{n+2}\Theta(y) = \Theta(0 \dots 000y) = \Theta(x^{\tilde{f}}).$$

□

4.2. Fibonacci sequence. From now on, we identify the semigroup F with $\Gamma(I)$ and use the notation F for both. Recall the definition of the following elements of F : $f_1 = s$, $f_2 = f$, and $f_n = f_{n-2}f_{n-1}$ for $n \geq 3$, and the Fibonacci numbers Φ_n defined by $\Phi_1 = \Phi_2 = 1$ and $\Phi_n = \Phi_{n-2} + \Phi_{n-1}$ for $n \geq 3$. We note that $\|f_n\| = \Phi_n$ for all $n \geq 1$. Set $\varphi = (1 + \sqrt{5})/2$; recall that $\Phi^n \approx \varphi^n/\sqrt{5}$.

Note that if n is even and ≥ 6 , then f_n starts with $fsfs$, while if n is odd and ≥ 5 then f_n starts with $sffs$. The following statements are easily proven by induction:

Lemma 4.2. (1) For all $k \leq n-2$ with $k \equiv n \pmod{2}$ we have $f_n \equiv f_k(f_{k+1}f_{k+3} \dots f_{n-1})$; in particular, f_k is a prefix of f_n .

(2) For all $k > 1$ we have $f_{k+1}^2 \equiv f_{k-1}f_{k+2}$.

4.3. Some relations. By (7), we have

$$(8) \quad \ll g_0, g_1 \gg \zeta \ll h_0, h_1 \gg \zeta = \ll g_0 h_0, g_1 h_0 \gg \zeta$$

for all $g_i, h_i \in F$, a calculation that we will use repeatedly; so by direct computation

Lemma 4.3. In F the following relations hold: $s^2 = e$; $f^3 = f$.

Proof. If $x = x_1 \dots x_n$, then $x^{s^2} = ((1 - x_1)x_2 \dots x_n)^s = x$, so $s^2 = e$. Then

$$\begin{aligned} \phi(f^2) &= \ll s, f \gg \zeta \ll s, f \gg \zeta = \ll s^2, fs \gg \zeta, \\ \phi(f^3) &= \ll s^2, fs \gg \zeta \ll s, f \gg \zeta = \ll s^3, fs^2 \gg \zeta = \ll s, f \gg \zeta = f. \end{aligned} \quad \square$$

4.4. Contraction. The proofs will ultimately all rely on some form of induction on the length of representations of semigroup elements as words over $\{s, f\}$.

Lemma 4.4. The semigroup F is contracting.

Proof. Consider $g \in F$, and write it as a word w of minimal length $w_1 \dots w_n$; write also $\phi(g) = \ll g_0, g_1 \gg \pi_g$. Then, by Lemma 4.3, there cannot be more than two f 's in a row in w , so every group of three letters $w_{3k+1}w_{3k+2}w_{3k+3}$ contains at least an s , which will contribute no letter to g_0 nor g_1 . The other two letters contribute at most one letter each to each of g_0 and g_1 . In total, $\|g_0\| \leq \frac{2}{3}(\|g\| + 2)$, and similarly for $\|g_1\|$. □

Note, in fact, that the contraction ratio η may be chosen as $1/\varphi$, with a little more care. This becomes apparent in the next result.

Lemma 4.5. The decomposition of f_n satisfies

$$\begin{aligned} \phi(f_1) &= \ll e, e \gg \sigma, & \phi(f_2) &= \ll s, f \gg \zeta, \\ \phi(f_3) &= \ll f, s \gg \zeta, & \phi(f_4) &= \ll f_3, f^2 \gg \zeta, \\ \phi(f_{2n}) &= \ll f_{2n-1}, f_4 f_6 \dots f_{2n-2} \gg \zeta \text{ if } n \geq 3, \\ \phi(f_{2n+1}) &= \ll f_{2n}, f_2(f_5 f_7 \dots f_{2n-1}) \gg \zeta \text{ if } n \geq 2. \end{aligned}$$

Proof. The first four claims may be checked directly; we note that if $\phi(g) = \ll g_0, g_1 \gg \zeta$, then $\phi(sg) = \ll g_1, g_0 \gg \zeta$. Next, by induction,

$$\begin{aligned}\phi(f_{2n}) &= \phi(f_{2n-2})\phi(f_{2n-1}) = \ll f_{2n-3}, f_4 f_6 \cdots f_{2n-4} \gg \zeta \ll f_{2n-2}, * \gg \zeta \\ &= \ll f_{2n-3} f_{2n-2}, f_4 f_6 \cdots f_{2n-4} f_{2n-2} \gg \zeta\end{aligned}$$

as claimed, where $*$ stands for an irrelevant element of F ; and

$$\begin{aligned}\phi(f_{2n+1}) &= \phi(f_{2n-1})\phi(f_{2n}) = \ll f_{2n-2}, f_2(f_5 \cdots f_{2n-3}) \gg \zeta \ll f_{2n-1}, * \gg \zeta \\ &= \ll f_{2n-2} f_{2n-1}, f_2(f_5 \cdots f_{2n-3} f_{2n-1}) \gg \zeta.\end{aligned}\quad \square$$

5. PROOFS

We start by finding relations in F ; we then show that they yield a simple normal form for elements of F .

5.1. Relations. Recall that for $n \geq 1$ we defined in (2,3) words r_n and r'_n over the alphabet $\{s, f\}$ by $r_1 = s^2$, $r'_1 = e$, and

$$\begin{aligned}r_n &= f_{n+1} f_n^2 \equiv f_{n-1} f_n^3 \quad (\equiv f_{n+1} f_{n-2} f_{n+1} \text{ if } n \geq 3), \\ r'_n &= f_{n\%2+1} (f_{n\%2+5} f_{n\%2+7} \cdots f_{n-1} f_{n+1}) f_n.\end{aligned}$$

Note that r'_n , for odd $n \geq 3$, can be obtained from r_n by removing the first four letters; indeed

$$\begin{aligned}(9) \quad f s f r'_n &\equiv f_4 f_1 \cdot f_2(f_6 \cdots f_{n+1}) f_n \equiv f_4 f_3(f_6 f_8 \cdots f_{n+1}) f_n \equiv f_4 f_5^2(f_8 \cdots f_{n+1}) f_n \\ &\equiv f_6 f_5(f_8 \cdots f_{n+1}) f_n \equiv \cdots \equiv f_{n-3} f_{n-2}^2 f_{n+1} f_n \equiv f_{n-1} f_n^3 \equiv r_n.\end{aligned}$$

Similarly, r'_n for even $n \geq 4$ can be obtained from r_n by removing the first three letters and replacing them by s ; indeed, writing $r'_n = s r''_n$,

$$\begin{aligned}(10) \quad s f f r''_n &\equiv f_3 f_2(f_5 \cdots f_{n+1}) f_n \equiv f_3 f_2(f_5 f_7 \cdots f_{n+1}) f_n \equiv f_3 f_4^2(f_7 \cdots f_{n+1}) f_n \\ &\equiv f_5 f_6^2(f_9 \cdots f_{n+1}) f_n \equiv \cdots \equiv f_{n-3} f_{n-2}^2 f_{n+1} f_n \equiv f_{n-1} f_n^3 \equiv r_n.\end{aligned}$$

Let e_n be the word obtained from f_n by deleting its first two symbols.

Lemma 5.1. *If $n \geq 4$, then we have $\phi(f_n) = \ll f_{n-1}, e_{n-1} \gg \zeta$.*

Proof. Assume first that n is even. By Lemma 4.2 we have $f_{n-1} = f_3 f_4 f_6 \cdots f_{n-2}$, so $e_{n-1} = f_4 f_6 \cdots f_{n-2}$, and the Lemma holds by Lemma 4.5. Similarly, if n is odd then $e_{n-1} = f_2 f_5 \cdots f_{n-2}$ by Lemma 4.2, and again the Lemma holds by Lemma 4.5. $\square$

Lemma 5.2. *In F the relations $r_n = r'_n$ hold for all $n \geq 1$.*

Proof. The cases $n \leq 2$ are covered in 4.3, since we have $r_2 \equiv s f^3 = s f \equiv r'_2$. Let us therefore assume $n \geq 3$. We follow the notation of Lemma 5.1. We have the decomposition

$$\phi(r_n) = \ll f_n f_{n-1}^2, e_n f_{n-1}^2 \gg \zeta = \ll r_{n-1}, f s r'_{n-1} \gg \zeta.$$

Suppose first that n is even. Then by (8) we have

$$\begin{aligned}\phi(r'_n) &= \phi(f_1(f_5 f_7 \cdots f_{n+1}) f_n) = \ll e_4, f_4 \gg \zeta \ll f_6, * \gg \zeta \cdots \ll f_n, * \gg \zeta \ll f_{n-1}, * \gg \zeta \\ &= \ll f_2(f_6 f_8 \cdots f_n) f_{n-1}, f_4 f_6 \cdots f_n f_{n-1} \gg \zeta = \ll r'_{n-1}, f s r'_{n-1} \gg \zeta,\end{aligned}$$

where $*$ stands for an element of F that is not relevant to the calculation. If n is odd, then similarly

$$\begin{aligned}\phi(r'_n) &= \phi(f_2 f_6 \cdots f_{n+1} f_n) = \ll f_1, f_2 \gg \zeta \ll f_5, * \gg \zeta \cdots \ll f_n, * \gg \zeta \ll f_{n-1}, * \gg \zeta \\ &= \ll f_1 f_5 \cdots f_n f_{n-1}, f_2 f_5 \cdots f_n f_{n-1} \gg \zeta = \ll r'_{n-1}, f s r'_{n-1} \gg \zeta.\end{aligned}$$

We have $r_{n-1} = r'_{n-1}$ by induction, so $r_n = r'_n$. $\square$

5.1.1. More relations. We will ultimately show that $\{r_n = r'_n\}$ is a complete set of relations for F ; however, we will first describe more relations, that are consequences of these but allow much faster simplifications.

Lemma 5.3. *Consider $a, b \in \mathbb{N}$ with $a \geq 3$ and $a \geq b + 2$. Then*

$$f_a f_b^2 = f_a.$$

Proof. We proceed by induction on b . Since $a \geq b + 2$, the word f_a ends with f_{b+2} ; it therefore suffices to show that $f_{b+2} f_b^2 = f_{b+2}$. We also note that the statement holds for $b \leq 2$, since then $f_3 f_1^2 = f_3$ and $f_4 f_2^2 = f_4$ by Lemma 4.3. Then for $b \geq 3$ we have

$$\begin{aligned}f_{b+2} f_b^2 &= f_b f_{b+1} f_b^2 = f_b r_b = f_b r'_b \\ &= f_b f_{1+b\%2} f_{5+b\%2} f_{7+b\%2} \cdots f_{b-1} f_{b+1} f_b \\ &= f_b f_{1+b\%2}^2 f_{2+b\%2} f_{4+b\%2} f_{7+b\%2} f_{9+b\%2} \cdots f_{b-1} f_{b+1} f_b \\ &= f_b f_{2+b\%2}^2 f_{3+b\%2} f_{7+b\%2} f_{9+b\%2} \cdots f_{b-1} f_{b+1} f_b \\ &= f_b f_{3+b\%2} f_{7+b\%2} f_{9+b\%2} \cdots f_{b-1} f_{b+1} f_b \\ &= \cdots = f_b f_{b-3} f_{b+1} f_b = f_b f_{b-3}^2 f_{b-2}^2 f_{b-1} f_b = f_b f_{b-1} f_b = f_{b+2}.\end{aligned} \quad \square$$

Lemma 5.4. *For all $n \geq 2$ we have*

$$r'_n = (f_1^2 f_2^2 f_3^2 \cdots f_{n-3}^2 f_{n-2}^2) f_{n+1}.$$

Proof. If $n = 2$ then $r'_2 = f_3$ and the Lemma holds. If $n \geq 3$, then

$$\begin{aligned}r'_n &= f_{1+n\%2} f_{5+n\%2} \cdots f_{n-1} f_{n+1} f_n \\ &= f_{1+n\%2}^2 f_{2+n\%2}^2 f_{3+n\%2} f_{7+n\%2} \cdots f_{n-1} f_{n+1} f_n \\ &= \cdots = f_{1+n\%2}^2 f_{2+n\%2}^2 f_{3+n\%2}^2 \cdots f_{n-3}^2 f_{n-2}^2 f_{n-1} f_n.\end{aligned} \quad \square$$

Lemma 5.5. *The equality $f_n^4 = f_n^2$ holds in F for $n \leq 4$, and $f_n^5 = f_n^3$ holds in F for all $n \geq 1$.*

Proof. The first statement follows from the definition of r_n . It suffices to check the second one for $n \geq 5$. Using Lemmata 5.4 and 5.3, we have

$$\begin{aligned}f_n^5 &\equiv f_n (f_{n-2} f_{n-1}) f_n^3 \equiv f_n f_{n-2} r_n = f_n f_{n-2} r'_n = f_n f_{n-2} f_1^2 f_2^2 f_3^2 \cdots f_{n-3}^2 f_{n-2}^2 f_{n+1} \\ &= \cdots = f_n f_{n-2} f_{n-3}^2 f_{n-2}^2 f_{n+1} = f_n r'_{n-3} f_{n-2}^2 f_{n+1} = f_n f_1^2 f_2^2 \cdots f_{n-5}^2 f_{n-2}^3 f_{n+1} \\ &= \cdots = f_n f_{n-2} f_{n+1} = f_n f_{n-2} f_{n-1} f_n = f_n^3.\end{aligned} \quad \square$$

Lemma 5.6. *For $n \geq 8$ we have*

$$\begin{aligned}f_{n-2} f_{n-3} f_{n-5} f_{n-7} \cdot f_{n-2} f_n &= f_{n-2} f_{n-3} f_{n-5} \cdot f_{n-3} f_n \\ &= f_{n-2} f_{n-3} \cdot f_{n-4} f_n = f_{n-2} \cdot f_{n-5} f_n = f_n.\end{aligned}$$

Proof. At first we prove by induction that for $n \geq 1$ we have

$$f_{n+1}f_n^2f_{n+2} = f_{n+3}.$$

The induction starts with two cases: for $n = 1$ we have $f_2f_1^2f_3 = f_2f_3 = f_4$, and for $n = 2$ we have $f_3f_2^2f_4 = f_3f_4 = f_5$. Using twice Lemma 5.4 and the induction hypothesis, we have for $n \geq 3$

$$\begin{aligned} f_{n+1}f_n^2f_{n+2} &= f_{n-1}f_n^3f_{n+1} \\ &= f_{n-1}f_{n-2}\underbrace{f_{n-1}f_n^3}_{r'_n}f_{n+1} = f_{n-1}f_{n-2}f_1^2f_2^2 \cdots f_{n-2}^2f_{n+1}^2 \\ &= f_{n-1}\underbrace{f_{n-2}f_{n-3}^2}_{r'_{n-3}}f_{n-2}^2f_{n+1}^2 = f_{n-1}f_1^2f_2^2 \cdots f_{n-5}^2f_{n-2}^3f_{n+1}^2 \\ &= f_{n-1}f_{n-2}^3f_{n+1}^2 = f_{n-1}f_{n-2}^2 \cdot f_{n-2}f_{n-1}f_n f_{n+1} \\ &= f_{n-1}f_{n-2}^2f_n \cdot f_{n+2} = f_{n+1}f_{n+2} = f_{n+3}. \end{aligned}$$

Using the arguments above, we carry out the following transformations

$$\begin{aligned} f_{n-2}f_{n-3}f_{n-5}f_{n-7} \cdot f_{n-2}f_n &= \\ &= f_{n-2}\underbrace{f_{n-3}f_{n-5}f_{n-7} \cdot f_{n-6}}_{f_{n-3}f_{n-5}^2 = f_{n-3}}f_{n-5}f_{n-3}f_n \\ &= f_{n-2}f_{n-3}f_{n-5} \cdot f_{n-3}f_n = f_{n-2}f_{n-3}f_{n-5} \cdot f_{n-5}f_{n-4}f_n \\ &= f_{n-2}f_{n-3} \cdot f_{n-4}f_n = f_{n-2}f_{n-3}f_{n-4}^2f_{n-3}f_{n-1} \\ &= f_{n-2} \cdot f_{n-5}f_{n-4}f_{n-3}f_{n-1} = f_{n-2} \cdot f_{n-5}f_n \\ &= f_{n-2}f_{n-3}^2f_{n-1} = f_n. \end{aligned} \quad \square$$

5.2. A normal form for F . We first show that every semigroup element can be written in the form of Theorem 2.2. We write $\mathcal{N}$ for the set of words described there.

The words in $\mathcal{N}$ are those words that can be written as a product $f_{i_1} \cdots f_{i_m} \cdots f_{i_n}$ where the sequence $i_1, \dots, i_m$ is increasing by steps of at least two, and the sequence $i_m, \dots, i_n$ is decreasing by steps of at least one.

An arbitrary word $w \in \{s, f\}^*$ is put into normal form by replacing subwords that are left-hand sides of the rules (N1)-(N10) by their right-hand side, until no such substitution is possible. For example, consider the possible reductions that can be performed on the word $f_4f_3^2f_4$:

$$\begin{array}{ccccc} f_4f_3^2f_4 & \xrightarrow{\text{N2: } f_3f_4 \rightarrow f_5} & f_4f_3f_5 & \xrightarrow{\text{N8: } p=1, a=3, q=2} & f_2f_5 & \xrightarrow{\text{N5: } f_2 \rightarrow f_1f_3} & f_1f_3f_5. \\ \downarrow \text{N3: } f_3^2 \rightarrow f_1f_4 & & \uparrow \text{N2: } f_1f_2 \rightarrow f_3 & & \uparrow \text{N3: } f_4^2 \rightarrow f_2f_5 & & \\ f_4f_1f_4^2 & \xrightarrow{\text{N3: } f_4^2 \rightarrow f_2f_5} & f_4f_1f_2f_5 & \xrightarrow{\text{N10: } p=3, a=1, q=3} & f_4^2 & & \end{array}$$

Lemma 5.7. *The rewriting system in Theorem 2.3 is terminating.*

Proof. We define a function $\eta = (\eta_1, \eta_2) : \{f_1, f_2, \dots\}^* \rightarrow \mathbb{N}^2$ with the lexicographic ordering, such that if $w \rightarrow w'$ is obtained by an elementary reduction then

$\eta(w) > \eta(w')$. Since $\mathbb{N}^2$ is well ordered this will prove that the rewriting system is terminating.

Consider therefore $w \in \{f_1, f_2, \dots\}^*$, say $w = f_{i_1} \dots f_{i_n}$. Define

$$\eta_1(w) = \begin{cases} \sum_{j=1}^n \Phi_{i_j} - 3 & \text{if } n \geq 2 \text{ and } i_1 = 1 \text{ and } i_2 \geq 3 \text{ is odd,} \\ \sum_{j=1}^n \Phi_{i_j} & \text{otherwise;} \end{cases}$$

$$\eta_2(w) = \sum_{j=1}^n (n+1-j)i_j.$$

We may then check that $\eta_1(w) > \eta_1(w')$ if w' is obtained from w using one of the rules (N1,4–5,8–10), while $\eta_1(w) = \eta_1(w')$ and $\eta_2(w) > \eta_2(w')$ if w' is obtained from w using one of the rules (N2–3,6–7).

For example, in rule (N9), only relation $r_{a+2} \rightarrow r'_{a+2}$ is used, and it shortens the length (as a word over the alphabet $\{s, f\}$) of w by 4; note that the length of w is precisely $\sum \Phi_{i_j}$. Write $w' = f_{i'_1} \dots f_{i'_{n'}}$; then

$$\eta_1(w) \geq \sum_{j=1}^n \Phi_{i_j} - 3 > \sum_{j=1}^n \Phi_{i_j} - 4 = \sum_{j=1}^{n'} \Phi_{i'_j} \geq \eta_1(w').$$

Consider, as another example, rule (N6). We have $\Phi_{a+1} + \Phi_a + \Phi_{a+3} = \Phi_{a+3} + \Phi_{a+2}$, so $\eta_1(w) = \eta_1(w') = \sum \Phi_{i_j}$. Say that the left-hand side of the rule appears at positions $n-j$, $n-j+1$ and $n-j+2$ in w ; then

$$\begin{aligned} \eta_2(w) - \eta_2(w') &\geq (j+1)(a+1) + ja + (j-1)(a+3) - j(a+3) - (j-1)(a+2) \\ &= (j+1)(a-1) + 1 > 0. \end{aligned}$$

All other rules are proven to decrease η in a similar fashion. $\square$

The following calculations will be useful in the proof of Lemma 5.9. Let $u = f_{i_1} \dots f_{i_m} \dots f_{i_n}$ be an arbitrary semigroup word written in normal form such that $n \geq 1$, and $i_1, i_n > 1$, and i_m is the maximal index. If $i_1 = 1$ or $i_n = 1$, the argument below should be applied to the word obtained from u by removing these instances of f_1 , adapting the statements accordingly. If $i_1 = 3$, let k be the largest integer such that $i_r = 2r+1$ for all $r < k$; if $i_1 = 4$, let k be the largest integer such that $i_r = 2r+2$ for all $r < k$; if $i_1 \geq 5$, then k need not be defined. Then, by Lemma 4.5, the element u affords the decomposition $\phi(u) = \ll u_0, u_1 \gg \zeta$, where

$$(11) \quad u_0 = \begin{cases} f_{i_1-1} \dots f_{i_m-1} \dots f_{i_n-1} & \text{if } i_1 > 3, \\ f_1 f_{2k-1} f_{i_k-1} \dots f_{i_n-1} & \text{if } i_1 = 3, \end{cases}$$

$$(12) \quad u_1 = \begin{cases} (f_1 f_3 \dots f_{i_1-2}) f_{i_2-1} \dots f_{i_n-1} & \text{if } i_1 \text{ is odd and } i_2 \neq i_1 - 1, \\ (f_4 f_6 \dots f_{i_1-1}) f_{i_3-1} \dots f_{i_n-1} & \text{if } i_1 \text{ is odd and } i_2 = i_1 - 1, \\ (f_4 f_6 \dots f_{i_1-2}) f_{i_2-1} \dots f_{i_n-1} & \text{if } i_1 \text{ is even } \geq 6 \text{ and } i_2 \neq i_1 - 1, \\ (f_1 f_3 \dots f_{i_1-1}) f_{i_3-1} \dots f_{i_n-1} & \text{if } i_1 \text{ is even } \geq 6 \text{ and } i_2 = i_1 - 1, \\ f_1 f_{2k-1} f_{2k-2} f_{i_k-1} \dots f_{i_n-1} & \text{if } i_1 = 4, i_k \leq 2k-2, \\ f_{i_2-1} f_{i_3-1} \dots f_{i_n-1} & \text{if } i_1 = 4, i_k = 2k-1 \text{ or } i_k \geq 2k+3. \end{cases}$$

The maximal index of u_1 is always less than i_m , but the maximal index of u_0 is equal to i_m if and only if $i_1 = 3$ and $i_k \leq 2k-2$.

As an example, consider $u = f_1 f_3 f_8$ and $v = f_3 f_5 f_7$. For u , we first compute $\phi(f_3 f_8) = \ll f_1 f_3 f_7, f_1 f_7 \gg \zeta$ using the first lines in (11) and (12) and $k = 2$; so

$$\phi(u) = \ll f_1 f_7, f_1 f_3 f_7 \gg \zeta. \quad \text{Similarly, } \phi(v) = \ll f_1 f_7, f_1 f_4 f_6 \gg \zeta,$$

using the first lines in (11) and (12) and $k = 4$. Note that $u_0 = v_0$ while $u \neq v$; this happened because one of u, v started with f_1 .

Lemma 5.8. *The equations $sfsf^2x = fx$ and $sfy = fsy$ have no solution in F .*

Proof. We will show that if the first equation had a solution x , then the second equation would have a solution y that is not longer than x ; and if the second equation had a solution y , then the first equation would have a solution x that is shorter than y .

Assume therefore that x is a solution to the first equation, and write $\phi(x) = \ll x_0, x_1 \gg \pi$. Then $\phi(sfsf^2x) = \ll f^2sx_0, sfsx_0 \gg \zeta\pi = \ll sx_0, fx_0 \gg \zeta\pi$, so in particular, by considering the second coördinate, $y = sx_0$ is a solution to the second equation, and is not longer than x .

Assume then that y is a solution to the second equation, and write $\phi(y) = \ll y_0, y_1 \gg \rho$. Then $\ll fy_0, sy_0 \gg \zeta\rho = \ll sy_1, fy_1 \gg \zeta\sigma\rho$, so $y_0 = sfy_1 = sfs(sy_1) = sfsfy_0$ and similarly $y_1 = sfsfy_1$. Clearly $y \notin \langle s \rangle$; so $y = s^i fz$ for some $i \in \mathbb{N}$ and $z \in F$. If i is odd then y_0 begins with f , while if i is even then y_1 begins with f . In all cases one of y_0, y_1 begins with f , say $y_i = fx$; then x is a solution to the first equation, and is shorter than y . $\square$

Lemma 5.9. *All elements of $\mathcal{N}$ are distinct in F .*

Proof. The elements e and f_1 are distinct since they act differently on X . In addition, they differ from all other elements of F since they are invertible.

Consider now two distinct normal forms

$$u = f_1^\delta f_{i_1} \cdots f_{i_m} \cdots f_{i_n}, \quad v = f_1^\epsilon f_{j_1} \cdots f_{j_p} \cdots f_{j_q},$$

with $i_m, j_p \geq 3$ and $i_m \geq j_p$.

If $i_n = 1$ and $j_q \neq 1$, then $0^u = 1$ while $0^v = 0$ so they are distinct; if $i_n \neq 1$ and $j_q = 1$ the same argument applies; if $i_n = j_q = 1$ then we may cancel both from the normal form and proceed. Let us therefore suppose $i_n, j_q > 1$.

We proceed by induction on the maximal index i_m , to prove that all normal forms with at most that maximal index are distinct in F .

The induction starts with the elements $f_3, f_3 f_2, f_1 f_3$ and $f_1 f_3 f_2$. They afford the following decompositions:

$$\begin{aligned} \phi(f_3) &= \ll f_1 f_3, f_1 \gg \zeta, & \phi(f_3 f_2) &= \ll f_1 f_3 f_1, e \gg \zeta, \\ \phi(f_1 f_3) &= \ll f_1, f_1 f_3 \gg \zeta, & \phi(f_1 f_3 f_2) &= \ll e, f_1 f_3 f_1 \gg \zeta. \end{aligned}$$

The first coördinates of these decompositions are different by the remarks above. Let us therefore suppose $i_m \geq 4$.

If $\delta = \epsilon = 1$: up to left-multiplying u and v by f_1 , we may replace δ and ϵ by 0 and proceed.

If $\delta = \epsilon = 0$: then

$$\begin{aligned} \phi(u) &= \ll f_{i_1-1} \cdots f_{i_m-1} \cdots f_{i_n-1}, u_1 \gg \zeta, \\ \phi(v) &= \ll f_{j_1-1} \cdots f_{j_p-1} \cdots f_{j_q-1}, v_1 \gg \zeta, \end{aligned}$$

where u_1 and v_1 are defined by (12).

If $i_1, j_1 > 3$, then the first coördinates of $\phi(u)$ and $\phi(v)$ are in normal form, and are distinct by assumption, so by induction they are distinct in F . Similarly, if $i_1 = j_1 = 3$ then the second coördinates equal $f_1 f_{i_2-1} \cdots f_{i_n-1}$ and $f_1 f_{j_2-1} \cdots f_{j_q-1}$, and they are in normal form, so by induction u and v are distinct in F .

We consider therefore the case $i_1 = 3, j_1 > 3$. Let k be maximal such that $i_r = 2r + 1$ for all $r < k$. Then the first coördinate of $\phi(v)$ is in normal form and starts by f_{j_1-1} while the first coördinate of $\phi(u)$, when put in normal form, is $f_1 f_{2k-1} f_{i_k-1} \cdots f_{i_n-1}$. By the next item or by the induction hypothesis these elements are distinct in F , whence u and v are distinct in F , too.

The case $i_1 = 3, j_1 > 3$ is considered similarly.

If $\delta \neq \epsilon$: by symmetry, we may assume $\delta = 0$ and $\epsilon = 1$. Then

$$\phi(v) = \ll e_{j_1-1} f_{j_2-1} \cdots f_{j_q-1}, f_{j_1-1} \cdots f_{j_p-1} \cdots f_{j_q-1} \gg \zeta.$$

Assume for contradiction that u, v are equal in F . Then, in F , we have

$$\begin{aligned} u_0 = f_{i_1-1} \cdots f_{i_n-1} &= v_1 = e_{j_1-1} f_{j_2-1} \cdots f_{j_q-1}, \\ u_1 = e_{i_1-1} f_{i_2-1} \cdots f_{i_n-1} &= v_0 = f_{j_1-1} \cdots f_{j_q-1}, \end{aligned}$$

so these words, when put in normal form, must be equal. It follows from Lemma 5.1 that $u_0 = s f u_1$ if i_1 is odd and $u_0 = f s u_1$ if i_1 is even. Similarly, we have $v_0 = s f v_1$ or $v_0 = f s v_1$, depending on the parity of j_1 .

We consider in turn all possibilities for the parity of i_1 and j_1 . If i_1 is even and j_1 is odd, then $u_0 = s f v_0 = s f f s v_1 = v_1$; it follows from (12) that when put in normal form, v_1 starts with f . Using Lemma 5.8, we reach the contradiction. The same argument holds, by symmetry, if i_1 is odd and j_1 is even.

If i_1 and j_1 are both even, then we find $(s f)^2 u_0 = u_0$; now by assumption u_0 is of the form $f_k \dots$ for an odd k , so its normal form starts with f ; this contradicts Lemma 5.8.

Finally, if i_1 and j_1 are both odd, then we find $(f s)^2 u_0 = u_0$, and u_0 is of the form $f_k \dots$ for an even k , so its normal form starts with s ; again this contradicts Lemma 5.8. □

Proof of Theorem 2.3. We saw in Lemma 5.7 that the rewriting system is terminating. On the other hand, consider a word $w \in \{f_1, f_2, \dots\}^*$ and two reduced words w', w'' that are gotten from w by applying two different sequences of elementary reductions. The reduced forms of the rewriting system correspond precisely to the normal form described in Theorem 2.2; and they define equal elements in F , so by Lemma 5.9 they are graphically equal. This proves that the rewriting system is confluent. □

Proof of Theorem 2.2. Every semigroup element can be represented by a word, and this word can be put in reduced form by applying the rules from Theorem 2.3. As noted above, this reduced form coincides with the normal form; therefore the natural map $\mathcal{N} \rightarrow F$ is onto. It is one-to-one by Lemma 5.9.

Finally, all the rules (N1-10) derive from the rules $s^2 = 1$ (N1) and $r_n \rightarrow r'_n$ for $n \geq 2$ (N4,8-10), and from graphical equalities (N2-3,5-7). Each of the rules $r_n \mapsto r'_n$ is strictly length-shortening, so $\mathcal{N}$ is precisely the set of irreducible words

with respect to the rules, and consists of minimal representative words. In other words, if there were a shorter representation w' of some $w \in \mathcal{N}$, then this shorter representation could be reduced to a word $w'' \in \mathcal{N}$ using the rules $r_n \mapsto r'_n$, still yielding a word shorter, and therefore different, from w . This contradicts Lemma 5.9.

Now $\|f_n\| = \Phi_n$, and the initial f_1 should cancel with the initial s of f_{i_1} if i_1 is odd; this explains the length formula. $\square$

Proof of Theorem 2.1. The relations $r'_n = r_n$ for $n \geq 1$ hold in F by Lemma 5.2. They are also sufficient; otherwise, there would be two distinct words w, w' that are unequal in the abstract semigroup $\langle s, f \mid r'_n = r_n \rangle$ but are equal in F ; by Theorem 2.2 the normal form representation of w and w' are equal; but then w and w' can be transformed into each other using only the relations $r_n = r'_n$, contradicting our initial assumption. $\square$

5.3. Growth.

Proof of Theorem 2.4. Consider a typical element in the ball of radius ℓ in F . It will have a minimal normal form as given by Theorem 2.2, whence $\Phi_{i_m} \leq \ell$. The maximal length of a normal form for that value of i_m is $\Phi_{i_m} + \Phi_{i_m-1} + 2\Phi_{i_m-2} + \Phi_{i_m-3} + 2\Phi_{i_m-4} + \dots < 2\Phi_{i_m+1}$, so it follows that $\ell < 2\Phi_{i_m+1}$. Now since $\Phi_i \approx \varphi^i/\sqrt{5}$, we obtain

$$\varphi^{i_m}/\sqrt{5} \leq \ell < 2/\sqrt{5}\varphi^{i_m+1},$$

so

$$\frac{\log(\sqrt{5}\ell/(2\phi))}{\log \phi} \leq i_m \leq \frac{\log(\sqrt{5}\ell)}{\log \phi}.$$

Given any value of i_m , there are 2^{i_m} choices of i_j to include in the normal form with $j > m$, and $2\Phi_{i_m-2}$ choices for the i_j to include with $j < m$: the 2 counts the possible values of i_1 , and the Φ_{i_m-2} counts the possible values of $(i_3, \dots, i_{m-2}) \in \{0, 1\}^{m-4}$ with no two consecutive 1's. It follows that the ball growth function $\gamma(\ell)$ is given by

$$\gamma(\ell) = \sum_{k=1}^{i_m} 2^k 2\Phi_{k-2} \approx \frac{2}{\sqrt{5}\varphi^2} \frac{(2\varphi)^{i_m} - 1}{2\varphi - 1},$$

which gives the upper and lower bounds

$$\frac{2}{\sqrt{5}\varphi^2(2\varphi - 1)} \left(\frac{\sqrt{5}\ell}{2\varphi} \right)^\alpha \leq \gamma(\ell) \leq \frac{2}{\sqrt{5}\varphi^2(2\varphi - 1)} (\sqrt{5}\ell)^\alpha,$$

with $\alpha = \log(2\varphi)/\log \varphi$. $\square$

5.4. Identities.

Proof of Theorem 2.5. More generally, we will show that all of the equalities $g(wg)^5 = g(wg)^3$, for all $w \in \{1, s, f, fs, f^2, f^2s\}$, hold in F . We prove this by considering simultaneously all these equalities, and proceed by induction on $\|g\|$.

The induction is easily checked if $\|g\| \leq 2$, by considering in turn the cases $g = 1, s, f, sf, fs, f^2$. For longer g , depending on whether g starts or ends with an s , we have $\phi(g) = \ll fh, sh \gg \zeta$ or $\ll sh, fh \gg \zeta$ or $\ll fh, sh \gg \zeta \sigma$ or $\ll sh, fh \gg \zeta \sigma$; for

simplicity, assume that we are in the first case. Then $\|h\| < \|g\|$ by Lemma 4.4, so the equalities above hold for h by induction. Then

$$\begin{aligned}
\phi(gg^5) &= \ll fh(fh)^5, sh(fh)^5 \gg \zeta = \phi(gg^3), \\
\phi(g(sg)^5) &= \ll fh(sh)^5, sh(sh)^5 \gg \zeta = \phi(g(sg)^3), \\
\phi(g(fg)^5) &= \ll fh(f^2h)^5, sh(f^2h)^5 \gg \zeta = \phi(g(fg)^3), \\
\phi(g(fsg)^5) &= \ll fh(fsh)^5, sh(fsh)^5 \gg \zeta = \phi(g(fsg)^3), \\
\phi(g(f^2g)^5) &= \ll fh(f^3h)^5, sh(f^3h)^5 \gg \zeta = \ll fh(fh)^5, sh(fh)^5 \gg \zeta = \phi(g(f^2g)^3), \\
\phi(g(f^2sg)^5) &= \ll fh(f^2sh)^5, sh(f^2sh)^5 \gg \zeta = \phi(g(f^2sg)^3). \quad \square
\end{aligned}$$

5.5. Ideal structure.

Proof of Theorem 2.6. Let us prove by induction on n that $z_n(u) = 0^n$ for any $u \in X^n$, and that $f_{n+2}f_{n+1} = z_n$ holds in W_n , i.e. that z_n and $f_{n+2}f_{n+1}$ are the left-hand zeroes over X^n . For $n = 1$ it follows from the definition of f_3 that $f_3 : X^1 \rightarrow \{0\}$, whence the relations $f_1f_3 = f_2f_3 = f_3$ and $f_3f_2 = f_3$ hold.

Choose $n \geq 2$. The element z_n has the following decomposition:

$$z_n = \ll f_2f_4 \cdots f_{n+1}, f_1f_4f_6 \cdots f_{n+1} \gg \zeta = \ll f_1f_3z_{n-1}, f_1z_{n-1} \gg \zeta$$

if n is odd, and

$$z_n = \ll f_3f_5 \cdots f_{n+1}, f_2f_2f_5f_7 \cdots f_{n+1} \gg \zeta = \ll z_{n-1}, f_1f_3f_1z_{n-1} \gg \zeta$$

if n is even. Clearly both coördinates of decompositions end with z_{n-1} , which is a left-hand zero over X^{n-1} by the induction hypothesis. As $z_n(x) = 0$ for $x \in X$, we have $z_n : X^n \rightarrow \{0^n\}$.

Similarly, the element $f_{n+2}f_{n+1}$ is written in the following way:

$$\begin{aligned}
f_{n+2}f_{n+1} &= \ll f_{n+1}f_n, f_1f_3 \cdots f_nf_n \gg \zeta \\
&= \ll f_{n+1}f_n, f_4 \cdots f_{n-1}f_{n+1} \gg \zeta = \ll f_{n+1}f_n, z_{n-1} \gg \zeta
\end{aligned}$$

if n is odd, and

$$\begin{aligned}
f_{n+2}f_{n+1} &= \ll f_{n+1}f_n, f_4f_6 \cdots f_nf_n \gg \zeta \\
&= \ll f_{n+1}f_n, f_1f_3 \cdots f_{n-1}f_{n+1} \gg \zeta = \ll f_{n+1}f_n, f_1z_{n-1} \gg \zeta
\end{aligned}$$

if n is even. By the induction hypothesis and the proof above for z_n , the element $f_{n+2}f_{n+1}$ is a left-hand zero over X^n .

As $z_n, f_{n+2}f_{n+1} : X^n \rightarrow \{0^n\}$, the relations $f_1z = f_2z = z$ and $f_{n+2}f_{n+1} = z$ hold in W_n . We show by induction on the maximal index of a semigroup element that W_n includes no elements with maximal index $> n + 2$. It follows from the definition of f_n that

$$f_{n+3} = f_{3-n\%2}f_{4-n\%2}f_{6-n\%2} \cdots f_{n+2} = f_{3-n\%2}z_n = z_n.$$

For any $k > n+3$ the equality $f_k = f_{k-2}f_{k-1}$ holds and by the induction hypothesis $f_k = f_{k-2}z_n = z_n$.

It is necessary to prove that the other semigroup elements define different transformations of X^n . Once again, we use induction on n . For $n = 1$ the four semigroup elements e, f_1, f_3 and f_3f_1 define different automatic transformations over X^1 .

Fix $n \geq 2$. Let us write

$$\begin{aligned} S_n &= \{g \in F \text{ with maximal index } n\}, \\ Z_n &= \{g \in S_{n+2} : g \text{ includes } z_n\}, \\ A_n &= \{g \in S_{n+2} : g \text{ includes neither } z_n \text{ nor } f_{n+2}f_{n+1}\}, \\ B_n &= S_{n+2} \setminus (Z_n \cup A_n). \end{aligned}$$

The semigroup W_n is constructed as $W_{n-1} \cup B_{n-1} \cup Z_n \cup A_n$. It follows from the proof above that elements with maximal index $n+2$ act as left-hand zeroes on X^{n-1} . Then elements of Z_n and A_n coincide with elements Z_{n-1} over X^{n-1} . On the other hand, let g be an arbitrary element from S_{n+2} . It follows from (12) that any input symbol decreases the maximal index of at least one coördinate in the decomposition of g . Hence, there exists the input word u of length $n-1$ such that it transforms g into the element with maximal index 3. As elements from S_3 can be distinguished over X^2 , elements from S_{n+2} cannot act as a left-hand zero over X^{n+1} . Therefore elements from Z_n can be distinguished from elements of $B_{n-1} \cup W_{n-1}$ over X^n .

Let g be an arbitrary element from A_n . As g does not include z_n and $f_{n+2}f_{n+1}$, the second (or the first, if g starts with f_1) coördinate of its decomposition either contains z_{n-2} and has maximal index n , or it does not include z_{n-1} and has maximal index $n+1$. In both cases the second coördinate does not act as a left-hand zero over X^{n-1} and therefore g isn't a left-hand zero over X^n . Consequently, $A_n \cap Z_n = \emptyset$, and all elements of W_n define different transformations of X^n .

Now we count the elements in W_n . An arbitrary element with maximal index $k \geq 3$ can be written as

$$f_1^{\alpha_1} \underbrace{f_3^{\alpha_3} \cdots f_{k-2}^{\alpha_{k-2}}}_{\Phi_{k-2} \text{ possibilities}} f_k \underbrace{f_{k-1}^{\beta_{k-1}} \cdots f_2^{\beta_2} f_1^{\beta_1}}_{2^{k-1} \text{ possibilities}},$$

where $\beta_j \in \{0, 1\}$ for all $j = 1, 2, \dots, k-1$, and $\alpha_i \alpha_{i+1} = 0$, $i = 3, \dots, k-2$. Therefore there are $2 \cdot \Phi_{k-2} \cdot 2^{k-1}$ elements with maximal index k . The semigroup W_n includes all elements with maximal index that is $< n+2$, semigroup elements of the form $z f_{k-1}^{\beta_{k-1}} \cdots f_2^{\beta_2} f_1^{\beta_1}$, and semigroup elements with maximal index $(n+2)$ that include neither z_n nor $f_{n+2}f_{n+1}$. Therefore,

$$\begin{aligned} \#W_n &= 2 + \sum_{k=3}^{n+1} 2^k \Phi_{k-2} + 2^n + \left(\frac{1}{2} 2^{n+2} \Phi_n - 2^{n+1} \right) \\ &= \sum_{k=1}^n 2^{k+2} \Phi_k - 2^n - 2^{n+1} \Phi_n + 2. \end{aligned} \quad \square$$

To prove Corollary 2.8, we begin by a Lemma:

Lemma 5.10. *In the semigroup F , we have*

$$(13) \quad f_n f_{n-1} = \begin{cases} f s f s f_{n+1} & \text{if } n \geq 4 \text{ is even,} \\ s f f s f_{n+1} & \text{if } n \geq 5 \text{ is odd;} \end{cases}$$

$$(14) \quad z_n = \begin{cases} f s f_{n+3} & \text{if } n \geq 2 \text{ is even,} \\ s f s f_{n+3} & \text{if } n \geq 3 \text{ is odd;} \end{cases}$$

$$(15) \quad f_n = \begin{cases} f s f s f_{n-1} f_{n-2} = f z_{n-3} & \text{if } n \geq 6 \text{ is even,} \\ s f f s f_{n-1} f_{n-2} = s f z_{n-3} & \text{if } n \geq 5 \text{ is odd;} \end{cases}$$

Proof. Induction on n , the basis of the induction being easily checked by applying the reduction algorithm. For example, take $n \geq 6$ even; then $f s f s f_{n+1} = f s f s f_{n-1} f_n = f_{n-2} f_{n-3} f_n$ by induction. Then $f_{n-2} f_{n-3} f_n = f_{n-2} f_{n-3} f_{n-2} f_{n-1} = f_{n-2} f_{n-1}^2 = f_n f_{n-1}$. All other cases are treated similarly. $\square$

Proof of Corollary 2.8. The proof follows from Theorem 2.6. Assume first that g includes z_{n-2} or $f_n f_{n-1}$. Then it acts as a left-hand zero on X^{n-2} , whence $\tau(g) \leq 2^{-(n-2)}$. On the other hand, g belongs to W_{n-1} and doesn't act as a left-hand zero on X^{n+1} ; therefore $\tau(g) = 2^{-(n-2)}$.

Now assume g includes neither z_{n-2} nor $f_n f_{n-1}$. Then g belongs to W_{n-2} , but acts as a left-hand zero on X^{n-3} . Therefore $g(X^*) = x_{\beta_1} x_{\beta_2} \dots x_{\beta_{n-3}} X^*$, whence $\tau(g) = 2^{-(n-3)}$.

Note next that the generator of the ideal I_n has trace 2^{-n} . By the proof above, every element of trace 2^{-n} is a multiple f_{n+3} , of $f_{n+2} f_{n+1}$, or of z_n . By Lemma 5.10, the ideals generated by these three elements coincide. $\square$

Proof of Theorem 2.9. Let g be an arbitrary element with maximal index $n \geq 3$, generating the ideal $I = FgF$. We show that there exist $\ell(g), r(g) \in F$ such that

$$\ell(g) \cdot g \cdot r(g) = \begin{cases} f_{n+1} & \text{if } g \text{ includes } z_{n-2} \text{ or } f_n f_{n-1} \text{ in its normal form,} \\ f_n & \text{otherwise.} \end{cases}$$

Therefore $I = FgF = Ff_k F = F_{2-k+3}$ by Corollary 2.8, where $k = n$ or $k = n+1$ depending on the cases above.

Let us assume that $g = f_n f_{n-1}^{\beta_{n-1}} \dots f_2^{\beta_2} f_1^{\beta_1}$, where $\beta_i \in \{0, 1\}$. Then we claim that for the element $g_2 = f_1^{\beta_1} f_2^{\beta_2} \dots f_{n-2}^{\beta_{n-2}}$ the equality

$$(16) \quad g \cdot g_2 = f_n f_{n-1}^{\beta_{n-1}} \dots f_2^{\beta_2} f_1^{\beta_1} \cdot f_1^{\beta_1} f_2^{\beta_2} \dots f_{n-2}^{\beta_{n-2}} = f_n f_{n-1}^{\beta_{n-1}}$$

holds. Applying $r_2 = r'_2$ or Lemma 5.3 if necessary,

$$f_n f_{n-1}^{\beta_{n-1}} \dots f_2^{\beta_2} f_1^{\beta_1} \cdot f_1^{\beta_1} f_2^{\beta_2} = f_n f_{n-1}^{\beta_{n-1}} \dots f_3^{\beta_3}.$$

If $\beta_i = 0, \beta_{i-1} = 1$ for some $i \geq 4$ then it follows from Lemma 5.3 that

$$f_n f_{n-1}^{\beta_{n-1}} \dots f_{i+1}^{\beta_{i+1}} f_i^0 f_{i-1} \cdot f_{i-1} f_i^0 = f_n f_{n-1}^{\beta_{n-1}} \dots f_{i+1}^{\beta_{i+1}}.$$

Otherwise, if $\beta_i = \beta_{i-1} = 1$ for some $i \geq 4$ then we have

$$f_n f_{n-1}^{\beta_{n-1}} \dots f_{i+1}^{\beta_{i+1}} f_i f_{i-1} \cdot f_{i-1} = f_n f_{n-1}^{\beta_{n-1}} \dots f_{i+1}^{\beta_{i+1}} f_i$$

by applying the relation

$$f_p f_m f_{m-1} \cdot f_{m-1} = f_p f_1^2 f_2^2 \dots f_{m-3}^2 f_m = f_p f_m$$

which holds for all $p > m \geq 2$, due to Lemmata 5.4 and 5.3.

Now let $g = f_3^{\alpha_3} \cdots f_{n-2}^{\alpha_{n-2}} f_n$ with $\alpha_i \alpha_{i+1} = 0$, $\alpha_i \in \{0, 1\}$, be such that g does not start with $f_3 f_5$ nor $f_4 f_6$.

At first we show that for any $m \geq 6$ and $k \leq m - 2$, with $(k, m) \neq (4, 6)$, there exists $w = w(k, m) \in F$ such that $w f_k f_m = f_m$. If $k = 3$ and $m = 6$ then $w = f_4 f_3 f_1$. If $m = 7$ then all cases of k are covered by the equalities

$$\begin{aligned} f_5 f_4 f_1 \cdot f_5 f_7 &= f_5 f_4 f_2 \cdot f_4 f_7 = f_5 f_4 \cdot f_3 f_7 \\ &= f_5 f_4 f_3^2 f_4 f_6 = f_5 f_4^2 f_6 = f_1 f_5 f_4 f_6 = f_1^2 f_5 f_4 f_5 = f_7. \end{aligned}$$

Using Lemma 5.6, the element $w(k, m)$ for $m \geq 8$ may be constructed as

$$w(k, m) = f_{m-2} f_{m-3} f_{m-5} f_{m-7} \cdot f_{m-3} f_{m-4} \cdots f_k f_{k-1}.$$

Indeed,

$$\begin{aligned} w(k, m) \cdot f_k f_m &= f_{m-2} f_{m-3} f_{m-5} f_{m-7} \cdot f_{m-3} f_{m-4} \cdots f_k f_{k-1} \cdot f_k f_m \\ &= f_{m-2} f_{m-3} f_{m-5} f_{m-7} \cdot f_{m-3} f_{m-4} \cdots f_k \cdot f_{k+1} f_m \\ &= \cdots = f_{m-2} f_{m-3} f_{m-5} f_{m-7} \cdot f_{m-2} f_m = f_m. \end{aligned}$$

Let $3 \leq i_1 < i_2 < \cdots < i_k < n - 1$ be the indices such that $\alpha_{i_j} = 1$, with the other $\alpha_j = 0$. Let us consider the element

$$g_1 = w(i_k, n) w(i_{k-1}, i_k) \cdots w(i_2, i_3) w(i_1, i_2).$$

It follows from Lemma 5.6 that

$$(17) \quad g_1 \cdot g = w(i_k, n) w(i_{k-1}, i_k) \cdots w(i_2, i_3) w(i_1, i_2) \cdot f_3^{\alpha_3} \cdots f_{n-2}^{\alpha_{n-2}} f_n = f_n.$$

Now we construct the elements $\ell(g)$ and $r(g)$. Choose

$$g = f_1^{\alpha_1} f_3^{\alpha_3} \cdots f_{n-2}^{\alpha_{n-2}} f_n f_{n-1}^{\beta_{n-1}} \cdots f_2^{\beta_2} f_1^{\beta_1}.$$

If g includes z_{n-2} then it has normal form $f_1^{\alpha_1} z_{n-2} f_{n-1}^{\beta_{n-1}} \cdots f_2^{\beta_2} f_1^{\beta_1}$, and we set $\ell(g) = f_{3-n \% 2} f_1^{\alpha_1}$ and $r(g) = g_2 f_{n-1}^{\beta_{n-1}}$. According to the arguments above, we have

$$\ell(g) \cdot g \cdot r(g) = f_{n+1} f_{n-1}^{2\beta_{n-1}} = f_{n+1}.$$

Otherwise g does not include z_{n-2} . Assume first that g starts neither with $f_3 f_5$ nor with $f_4 f_6$. We set $\ell(g) = g_1 f_1^{\alpha_1}$ and $r(g) = g_2$. It follows from (16) and the arguments above that

$$\ell(g) \cdot g \cdot r(g) = f_n f_{n-1}^{\beta_{n-1}}.$$

If $\beta_{n-1} = 0$, then we are done. If $\beta_{n-1} = 1$ it follows from Lemma 5.10 that it suffices to take $\ell(g) = f s f s \cdot g_1 f_1^{\alpha_1}$ if n is even, and $\ell(g) = s f f s \cdot g_1 f_1^{\alpha_1}$ if n is odd; then $\ell(g) \cdot g \cdot r(g) = f_{n+1}$.

The last case is when g starts with $f_1^{\alpha_1} f_3 f_5 \cdots f_m$ or $f_1^{\alpha_1} f_4 f_6 \cdots f_m$, with $6 \leq m < n - 2$. We may consider respectively the elements $f_2 f_1^{\alpha_1} g$ or $f_3 f_1^{\alpha_1} g$, which satisfy the conditions of the previous paragraph. $\square$

REFERENCES

- [1] Alexander G. Abercrombie, *Subgroups and subrings of profinite rings*, Math. Proc. Cambridge Philos. Soc. **116** (1994), no. 2, 209–222.
- [2] L. Bartholdi, I. I. Reznikov, and V. I. Sushchansky, *The smallest Mealy automaton of intermediate growth*, J. Algebra **295** (2006), no. 2, 387–414, available at [arXiv:math.GR/0407312](https://arxiv.org/abs/math.GR/0407312). MR **2194959** (2006i:68060)
- [3] A. Ya. Belov and I. A. Ivanov, *Construction of semigroups with some exotic properties*, Acta Appl. Math. **85** (2005), no. 1-3, 49–56. MR **2128898** (2005m:20138)

- [4] ———, *Construction of semigroups with some exotic properties*, Comm. Algebra **31** (2003), no. 2, 673–696. MR **1968920** (2004a:20064)
- [5] David B. A. Epstein, James W. Cannon, Derek F. Holt, Silvio V. F. Levy, Michael S. Paterson, and William P. Thurston, *Word processing in groups*, Jones and Bartlett Publishers, 1992.
- [6] Israel M. Gel'fand and Alexander A. Kirillov, *On fields connected with the enveloping algebras of Lie algebras*, Dokl. Akad. Nauk SSSR **167** (1966), 503–505.
- [7] Rostislav I. Grigorchuk, *On Burnside's problem on periodic groups*, Функционал. Анал. и Приложен. **14** (1980), no. 1, 53–54. English translation: Functional Anal. Appl. **14** (1980), 41–43.
- [8] Günter R. Krause and Thomas H. Lenagan, *Growth of algebras and Gelfand-Kirillov dimension*, Revised edition, Graduate Studies in Mathematics, vol. 22, American Mathematical Society, Providence, RI, 2000.
- [9] I. I. Reznikov and V. I. Sushchansky, *On the 3-state Mealy automata over an m -symbol alphabet of growth order $[n^{\log n/2 \log m}]$* , J. Algebra **304** (2006), 712–754.
- [10] Lev M. Shneerson, *Relatively free semigroups of intermediate growth*, J. Algebra **235** (2001), no. 2, 484–546.
- [11] Robert B. Warfield Jr., *The Gelfand-Kirillov dimension of a tensor product*, Math. Z. **185** (1984), no. 4, 441–447. MR **733766** (85f:17006)